

GRANULEX

INDICATIONS:

Varicose Ulcers
Decubital Ulcers

Debride Eschar
Sunburn

Dehiscent Wounds
Wound Healing

GRANULEX will relieve pain and promote healing; debrides eschar and necrotic tissue physiologically; stimulates vascular bed; improves epithelization by reducing premature epithelial desiccation and cornification; reduces odor from malodorous necrotic wounds.

DIRECTIONS: Shake well before spraying. Hold upright and approximately 12 inches from the area to be treated. Press valve and coat wound rapidly. Wound may be left unbandaged or apply a wet dressing. Apply twice daily or as often as necessary. To remove, wash gently with water. When applied to a sensitive area, a temporary stinging sensation may be noted.

WARNING: Do not use on fresh arterial clots. Avoid spraying in eyes. Flammable, do not expose to fire or open flame. Contents under pressure. Do not puncture or incinerate. Do not store at temperature above 120°F. Keep out of reach of children. Use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

301.2

Contents: **Net weight 2 oz. (56.7g)**
Inert propellant 25% of total content.



NDC No. 51079-621-81



SHAKE WELL BEFORE EACH USAGE
Hold Upright to Spray

The nozzle of the canister is designed for large volume delivery of medication, therefore, proper coverage of the wound site is achieved quickly.

PHYSIOLOGICAL DEBRIDEMENT
CAPILLARY BED STIMULANT
PROMOTES GRANULATION

Each gram contains:

Trypsin, USP 0.12 mg
Balsam Peru 87.0 mg
Castor Oil, USP 788.0 mg

With an emulsifier and propellants (Water Dispersible)

 **UDL Laboratories, Inc.**
Rockford, IL 61103

Rx only

Net Wt.
2 oz. (56.7g)